

AD-A169 968


A STRUCTURE THEOREM ON BIVARIATE POSITIVE QUADRANT  
DEPENDENT DISTRIBUTION. (U) PITTSBURGH UNIV PA CENTER  
FOR MULTIVARIATE ANALYSIS M. B. RAO ET AL. DEC 85


1/1

UNCLASSIFIED

TR-85-48 AFOSR-TR-86-0344 F49620-85-C-0000 F/G 12/1

NL





AD-A169 968



A STRUCTURE THEOREM ON BIVARIATE POSITIVE  
QUADRANT DEPENDENT DISTRIBUTIONS AND TESTS FOR  
INDEPENDENCE IN TWO-WAY CONTINGENCY TABLES\*

by

M. Bhaskara Rao\*\*  
P.R. Krishnaiah  
K. Subramanyam

**Center for Multivariate Analysis**  
**University of Pittsburgh**

ATC FILE COPY



Approved for public release;  
distribution unlimited.

A STRUCTURE THEOREM ON BIVARIATE POSITIVE  
QUADRANT DEPENDENT DISTRIBUTIONS AND TESTS FOR  
INDEPENDENCE IN TWO-WAY CONTINGENCY TABLES\*

by

M. Bhaskara Rao\*\*  
P.R. Krishnaiah  
K. Subramanyam

December 1985

Technical Report No. 85-48

DTIC  
ELECTE  
S JUL 23 1986 D  
B

Center for Multivariate Analysis  
Fifth Floor, Thackeray Hall  
University of Pittsburgh  
Pittsburgh, PA 15260

DISTRIBUTION STATEMENT A

Approved for public release  
Distribution Unlimited

\*This work was supported by the Air Force Office of Scientific Research under Contract F49620-85-C-0008. The U.S. Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon.

\*\*At present with Department of Probability and Statistics, University of Sheffield.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)  
NOTICE OF TRANSMITTAL TO DTIC  
This technical report has been reviewed and is  
approved for public release in accordance with AFM 190-12.  
Distribution is unlimited.  
MATTHEW J. KEEPER  
Chief, Technical Information Division

A STRUCTURE THEOREM ON BIVARIATE POSITIVE  
QUADRANT DEPENDENT DISTRIBUTIONS AND TESTS FOR  
INDEPENDENCE IN TWO-WAY CONTINGENCY TABLES\*

by

M. Bhaskara Rao\*\*  
P.R. Krishnaiah  
K. Subramanyam

ABSTRACT

In this paper, the set of all bivariate positive quadrant dependent distributions with fixed marginals is shown to be compact and convex. Extreme points of this convex set are enumerated in some specific examples. Applications are given in testing the hypothesis of independence against strict positive quadrant dependence in the context of ordinal contingency tables. Various procedures based upon certain functions of the eigenvalues of a random matrix are also proposed for testing for independence in two-way contingency table. The performance of some tests one of which is based on eigenvalues of a random matrix is compared.

AMS 1980 Subject Classifications: Primary 62H05; Secondary 62H17, 62H15

Key words and phrases: Asymptotic distributions, compact set, contingency tables, convex set, eigenvalues, extreme points, gamma ratio, hypothesis of independence, positive quadrant dependent distributions, power function.

\*This work was supported by the Air Force Office of Scientific Research under Contract F49620-85-C-0008. The U.S. Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon.

\*\*At present with Department of Probability and Statistics, University of Sheffield.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER <b>AFOSR-TR- 86 - 0344</b>	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A structure theorem on bivariate positive quadrant dependent distributions and tests for independence in two-way contingency tables.		5. TYPE OF REPORT & PERIOD COVERED Technical - December 1985
		6. PERFORMING ORG. REPORT NUMBER 85-48
7. AUTHOR(s) M. Bhaskara Rao, P.R. Krishnaiah, K. Subramanyam.		8. CONTRACT OR GRANT NUMBER(s) F49620-85-C-0008 (Air Force)
9. PERFORMING ORGANIZATION NAME AND ADDRESS Center for Multivariate Analysis 515 Thackeray Hall University of Pittsburgh, Pittsburgh, PA 15260		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102 F 2304/AS
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research Department of the Air Force Bolling Air Force DC 20332		12. REPORT DATE December 1985
		13. NUMBER OF PAGES 60
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		16a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Asymptotic distributions, compact set, contingency tables, convex set, eigenvalues, extreme points, gamma ratio, hypothesis of independence, positive quadrant dependent distributions, power function.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In this paper, the set of all bivariate positive quadrant dependent distributions with fixed marginals is shown to be compact and convex. Extreme points of this convex set are enumerated in some specific examples. Applications are given in testing the hypothesis of independence against strict positive quadrant dependence in the context of ordinal contingency tables. Various procedures based upon certain functions of the eigenvalues of a random matrix are also proposed for testing for independence in two-way contingency table. The		

DD FORM 1 JAN 73 1473

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)


performance of some tests one of which is based on eigenvalues of a random matrix is compared.

Unclassified

## 1. INTRODUCTION

Cross-classified data having ordered categories arise in many investigations conducted by medical, physical, natural and social scientists. Statistical methods have been developed and continue to evolve to analyze such data. Many of these methods are tailored to answer specific questions and issues raised. For reviews of the literature in this area, the reader is referred to Agresti [1] and Goodman [6].

We begin with a general description of a problem tackled in this paper. Let  $\mathcal{B}$  be the Borel  $\sigma$  - field on the real line,  $\mathcal{R}$  and  $\mathcal{B} \times \mathcal{B}$  the product  $\sigma$  - field on  $\mathcal{R} \times \mathcal{R}$ . Let  $\mu$  be a probability measure on  $\mathcal{B} \times \mathcal{B}$  and  $\mu_1$  and  $\mu_2$  the corresponding marginal probability measures on  $\mathcal{B}$ , i.e.,  $\mu_1(B) = \mu(B \times \mathcal{R})$  and  $\mu_2(B) = \mu(\mathcal{R} \times B)$  for every  $B$  in  $\mathcal{B}$ . Following Lehmann [14],  $\mu$  is said to be a positive quadrant





dependent if

$$\mu\{[c,\infty) \times [d,\infty)\} \geq \mu_1\{[c,\infty)\} \mu_2\{[d,\infty)\}$$

for every  $c, d$  in  $R$ . In the jargon of random variables, the above notion can be rephrased as follows. Let  $X$  and  $Y$  be two random variables with some joint probability distribution function  $F$ .  $X$  and  $Y$  are said to be positive quadrant dependent if

$$P\{X \geq c, Y \geq d\} \geq P\{X \geq c\} P\{Y \geq d\}$$

for all  $c, d$  in  $R$ . For various properties of positive quadrant dependence, see Lehmann [14] or Eaton [3]. In this paper, we look at the notion of positive quadrant dependence from a global point of view. Let  $M_{PQD}$  denote the set of all positive quadrant dependent probability measures  $\mu$  on  $\mathcal{B} \times \mathcal{B}$ . It is natural to think along the following lines. If  $M_{PQD}$  is a convex set and compact in some decent topology, then the set of extreme points of  $M_{PQD}$  will be non-empty. See Phelps [20]. Moreover, every member of  $M_{PQD}$  can be expressed as a mixture (in some sense) of extreme points of  $M_{PQD}$ . There are certain properties of distributions which are preserved under mixtures. Under these circumstances, it suffices to examine extensively the extreme points so as to make comments on the members of  $M_{PQD}$ . But this line of reasoning fails since  $M_{PQD}$  is not a convex set as the following example demonstrates.

Let  $\mu$  be a probability measure on  $\mathcal{B} \times \mathcal{B}$  with support contained in  $\{(1,1), (1,2), (2,1), (2,2)\}$ . Such a probability measure can be written as

$$\mu = \begin{array}{c|cc|c} & 1 & 2 & \\ \hline 1 & p_{11} & p_{12} & p_1 \\ 2 & p_{21} & p_{22} & p_2 \\ \hline & q_1 & q_2 & 1 \end{array}$$

where  $p_{ij} = \mu(\{(i,j)\})$ ,  $i = 1,2$ ;  $j = 1,2$ ;  $p_i = \mu_1(\{i\})$ ,  $i = 1,2$  and  $q_j = \mu_2(\{j\})$ ,  $j = 1,2$ . Then  $\mu \in M$  if and only if  $p_2 q_2 \leq p_{22} \leq p_2 \wedge q_2$ , where  $p_2 \wedge q_2$  denotes the minimum of  $p_2$  and  $q_2$ . For the desired example, let  $\mu$  and  $\eta$  be the probability measures with the same support  $\{(1,1), (1,2), (2,1), (2,2)\}$  given by

$$\mu = \begin{array}{c|cc|c} & 1 & 2 & \\ \hline 1 & \frac{1}{4} & \frac{1}{4} & \frac{1}{2} \\ 2 & \frac{1}{4} & \frac{1}{4} & \frac{1}{2} \\ \hline & \frac{1}{2} & \frac{1}{2} & 1 \end{array}$$

$$\eta = \begin{array}{c|cc|c} & 1 & 2 & \\ \hline 1 & \frac{2}{9} & \frac{1}{9} & \frac{1}{3} \\ 2 & \frac{4}{9} & \frac{2}{9} & \frac{2}{3} \\ \hline & \frac{2}{3} & \frac{1}{3} & 1 \end{array}$$

$\mu$  and  $\eta$  are positive quadrant dependent but  $\frac{1}{2}\mu + \frac{1}{2}\eta$  is not.

We can identify some natural subsets of  $M_{PQD}$  as convex sets. Let  $\lambda$  and  $\nu$  be two probability measures on  $\mathcal{B}$ . Let  $M_{PQD}(\lambda, \nu)$  be the collection of all probability measures  $\mu$  in  $M_{PQD}$  such that  $\mu_1 = \lambda$  and  $\mu_2 = \nu$ , i.e.,

$$M_{PQD}(\lambda, \nu) = \{\mu \in M_{PQD} ; \mu_1 = \lambda \text{ and } \mu_2 = \nu\}.$$

In Section 2, we show that  $M_{PQD}(\lambda, \nu)$  is a compact convex set in the weak topology on the space of all probability measures  $M$  on  $\mathcal{B} \times \mathcal{B}$ . Using this result, one obtains a decomposition of  $M_{PQD}$  as

$$M_{PQD} = \bigcup_{\lambda} \bigcup_{\nu} M_{PQD}(\lambda, \nu),$$

where the union is taken over all probability measures  $\lambda, \nu$  on  $\mathcal{B}$ .

In Section 3, we concentrate on the case when both  $\lambda$  and  $\nu$  have finite support. We describe a method of enumerating all extreme points of  $M_{PQD}(\lambda, \nu)$  with the help of some examples. In Section 4, using the structure of  $M_{PQD}(\lambda, \nu)$ , we compare the performance of some tests for testing independence against strict positive quadrant dependence.

## 2. Main Results

In this section, we show that for any two probability measures  $\lambda$  and  $\nu$  on  $\mathcal{B}$ ,  $M_{PQD}(\lambda, \nu)$  is compact and convex.

We need the following definitions and results in this connection.

Let  $(X, d)$  be a Polish space, i.e., a complete separable metric space. Let  $\mathcal{B}_X$  be the Borel  $\sigma$ -field on  $X$  and  $M_X$  the space of all probability measures on  $\mathcal{B}_X$ .  $M_X$  is equipped with weak topology.

Definition 1. A subset  $S$  of  $M_X$  is said to be uniformly tight if for every  $\epsilon > 0$  there exists a compact subset  $C$  of  $X$  such that

$$\mu(C) > 1 - \epsilon \quad \text{for every } \mu \text{ in } S.$$

The following is known as Prohorov's theorem.

Proposition 2 A subset  $S$  of  $M_X$  is relatively compact if and only if  $S$  is uniformly tight.  $S$  is compact if and only if  $S$  is closed and uniformly tight.

Proof. See Billingsley [ , Theorems 6.1 and 6.2, p.37 ].

Theorem 3 Let  $M(\lambda, \nu)$  be the collection of all probability measures  $\mu$  on  $\mathcal{B} \times \mathcal{B}$  such that  $\mu_1 = \lambda$  and  $\mu_2 = \nu$ . Then  $M(\lambda, \nu)$  is compact.

Proof. It is obvious that  $M(\lambda, \nu)$  is a closed subset of  $M$ , the space of all probability measures on  $\mathcal{B} \times \mathcal{B}$ . We show that  $M(\lambda, \nu)$  is uniformly tight. Let  $\epsilon > 0$ . There exist compact subsets  $C_1$  and  $C_2$  of  $R$  such that  $\lambda(C_1^c) < \epsilon/2$  and  $\nu(C_2^c) < \epsilon/2$ .  $C_1 \times C_2$  is a compact subset of  $R \times R$ . Let  $\mu \in M(\lambda, \nu)$ . Then

$$\begin{aligned}
 \mu[(C_1 \times C_2)^c] &\leq \mu(C_1^c \times R \cup R \times C_2^c) \\
 &\leq \mu(C_1^c \times R) + \mu(R \times C_2^c) \\
 &= \mu_1(C_1^c) + \mu_2(C_2^c) \\
 &= \lambda(C_1^c) + \nu(C_2^c) \\
 &< \varepsilon.
 \end{aligned}$$

This completes the proof in view of Proposition 2.

The following result is the main result of this section.

Theorem 4 For any given probability measures  $\lambda$  and  $\nu$  on  $\mathcal{B}$ ,  $M_{PQD}(\lambda, \nu)$  is compact and convex.

Proof.  $M_{PQD}(\lambda, \nu)$  is a closed subset of  $M(\lambda, \nu)$  follows from the following observation. Let  $\mu^n, n \geq 1$  be a sequence in  $M_{PQD}(\lambda, \nu)$  converging weakly to a  $\mu$  in  $M(\lambda, \nu)$ . Then for any  $c, d$  in  $R$ , (See Billingsley [2, p. 11]),

$$\begin{aligned}
 \mu\{[c, \infty) \times [d, \infty)\} &\geq \limsup_{n \rightarrow \infty} \mu^n\{[c, \infty) \times [d, \infty)\} \\
 &\geq \lambda\{[c, \infty)\} \nu\{[d, \infty)\}.
 \end{aligned}$$

Hence  $\mu \in M_{PQD}(\lambda, \nu)$ . This implies that  $M_{PQD}(\lambda, \nu)$  is compact. We now show that  $M_{PQD}(\lambda, \nu)$  is convex. Let  $\mu, \nu \in M_{PQD}(\lambda, \nu)$  and  $0 \leq \alpha \leq 1$ . Then for any  $c, d$  in  $R$ ,

$$\begin{aligned}
 & (\alpha\mu + (1-\alpha)\eta)\{[c,\infty) \times [d,\infty)\} - \lambda\{[c,\infty)\} \vee \{[d,\infty)\} \\
 &= \alpha\mu\{[c,\infty) \times [d,\infty)\} + (1-\alpha)\eta\{[c,\infty) \times [d,\infty)\} - \\
 &\quad \lambda\{[c,\infty)\} \vee \{[d,\infty)\} \\
 &\geq \alpha\lambda\{[c,\infty)\} \vee \{[d,\infty)\} + (1-\alpha)\lambda\{[c,\infty)\} \vee \{[d,\infty)\} \\
 &\quad - \lambda\{[c,\infty)\} \vee \{[d,\infty)\} = 0.
 \end{aligned}$$

Consequently,  $\alpha\mu + (1-\alpha)\eta \in M_{PQD}(\lambda, \nu)$ . This completes the proof.

### 3. Extreme Points

In this section, we assume that the support of  $\lambda$  is  $\{1,2,\dots,m\}$  and that of  $\nu$  is  $\{1,2,\dots,n\}$ . Let  $p_i = \mu(\{i\})$ ,  $i = 1,2,\dots,m$  and  $q_j = \nu(\{j\})$ ,  $j = 1,2,\dots,n$ . In this case, we use the suggestive notation  $M_{PQD}(p_1, p_2, \dots, p_m; q_1, q_2, \dots, q_n)$  for  $M_{PQD}(\lambda, \nu)$ . Any  $\mu$  in  $M_{PQD}(\lambda, \nu)$  can be written in the following form

$p_{11}$	$p_{12}$	$\dots$	$p_{1n}$	$p_1$
$p_{21}$	$p_{22}$	$\dots$	$p_{2n}$	$p_2$
$\dots$	$\dots$	$\dots$		$\vdots$
$p_{m1}$	$p_{m2}$	$\dots$	$p_{mn}$	$p_m$
$q_1$	$q_2$	$\dots$	$q_n$	1

where  $p_{ij} = \mu(\{(i,j)\})$ ,  $i = 1,2,\dots,m$ ;  $j = 1,2,\dots,n$ . In other words,  $M_{PQD}(p_1, p_2, \dots, p_m; q_1, q_2, \dots, q_n)$  is the collection of all matrices  $(p_{ij})$  of order  $m \times n$  such that each  $p_{ij} \geq 0$ , row sums  $p_1, p_2, \dots, p_m$ , column sums  $q_1, q_2, \dots, q_n$  and the joint distribution is positive quadrant dependent. The compact convex set  $M_{PQD}(p_1, p_2, \dots, p_m; q_1, q_2, \dots, q_n)$  has a finite number of extreme points. We now describe a method of enumerating the extreme points in some special cases for illustration from which the general technique can easily be perceived. As we shall see shortly in Section 4, the knowledge of extreme points has considerable bearing on the power of tests of independence against strict positive quadrant dependence.

Example 1.  $m = 2$  and  $n = 2$ .

Let  $p_1, p_2; q_1, q_2$  be specified such that each of  $p_1, p_2, q_1, q_2$  is positive and  $p_1 + p_2 = 1 = q_1 + q_2$ . It can be easily verified that if a matrix

$$\begin{bmatrix} p_{11} & p_{12} \\ p_{21} & p_{22} \end{bmatrix}$$

with non-negative entries, row sums  $p_1, p_2$  and column sums  $q_1, q_2$  belongs to  $M_{PQD}(p_1, p_2; q_1, q_2)$ , then

$$p_2 q_2 \leq p_{22} \leq p_2 \wedge q_2.$$

Conversely, if the number  $p_{22}$  satisfies the above inequality, then the matrix

$$\begin{bmatrix} p_1 - q_2 + p_{22} & q_2 - p_{22} \\ p_2 - p_{22} & p_{22} \end{bmatrix}$$

belongs to  $M_{PQD}(p_1, p_2; q_1, q_2)$ . There are only two extreme points of  $M_{PQD}(p_1, p_2; q_1, q_2)$ . These are given by

$$\begin{bmatrix} p_1 q_1 & p_1 q_2 \\ p_2 q_1 & p_2 q_2 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} q_1 & q_2 - p_2 \\ 0 & p_2 \end{bmatrix} \quad \text{if} \quad p_2 \wedge q_2 = p_2,$$

$$\begin{bmatrix} p_1 q_1 & p_1 q_2 \\ p_2 q_1 & p_2 q_2 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} p_1 & 0 \\ p_2 - q_2 & q_2 \end{bmatrix} \quad \text{if} \quad p_2 \wedge q_2 = q_2.$$

Every member of  $M_{PQD}(p_1, p_2; q_1, q_2)$  is a convex combination of these two extreme points.



Example 2  $m = 2$  and  $n = 3$ .

Let  $p_1, p_2, q_1, q_2, q_3$  be five positive numbers satisfying  $p_1 + p_2 = 1 = q_1 + q_2 + q_3$ . If the matrix  $(p_{ij})$  belongs to  $M_{PQD}(p_1, p_2; q_1, q_2, q_3)$ , then

$$p_2 q_3 \leq p_{23} \leq p_2 \wedge q_3 \quad \text{and}$$

$$(p_2 q_2 + p_2 q_3) \vee p_{23} \leq p_{22} + p_{23} \leq p_2 \wedge (q_2 + p_{23}),$$

where  $a \vee b$  indicates the maximum of the numbers  $a$  and  $b$ . Conversely, if  $p_{22}$  and  $p_{23}$  are two numbers satisfying the above inequalities, then the matrix


$$\begin{bmatrix} q_1 - p_2 + p_{22} + p_{23} & q_2 - p_{22} & q_3 - p_{23} \\ p_2 - p_{22} - p_{23} & p_{22} & p_{23} \end{bmatrix}$$

belongs to  $M_{PQD}(p_1, p_2; q_1, q_2, q_3)$ . The impact of this observation is that the numbers  $p_{22}$  and  $p_{23}$  in the matrix  $(p_{ij})$  determine whether the matrix  $(p_{ij})$  belongs to  $M_{PQD}(p_1, p_2; q_1, q_2, q_3)$  or not. These two inequalities determine a simplex in the  $p_{22} - p_{23}$  plane. As a simple illustration, take  $p_1 = p_2 = \frac{1}{2}$ ;  $q_1 = q_2 = q_3 = \frac{1}{3}$ . The determining inequalities are

$$\frac{1}{6} \leq p_{23} \leq \frac{1}{3} \quad \text{and}$$

$$\frac{1}{3} = p_{23} \vee \frac{1}{3} \leq p_{22} + p_{23} \leq \frac{1}{2} \wedge (\frac{1}{3} + p_{23}) = \frac{1}{2}.$$

These inequalities determine the following simplex in the  $p_{22} - p_{23}$  plane.



There are four extreme points of the set  $M_{PQD}(1/2, 1/2; 1/3, 1/3, 1/3)$  given by

$$\begin{bmatrix} 1/6 & 1/6 & 1/6 \\ 1/6 & 1/6 & 1/6 \end{bmatrix}, \quad \begin{bmatrix} 1/6 & 1/3 & 0 \\ 1/6 & 0 & 1/3 \end{bmatrix},$$

$$\begin{bmatrix} 1/3 & 1/6 & 0 \\ 0 & 1/6 & 1/3 \end{bmatrix}, \quad \begin{bmatrix} 1/3 & 0 & 1/6 \\ 0 & 1/3 & 1/6 \end{bmatrix}$$

corresponding to the four points  $P_1, P_2, P_3, P_4$  respectively. Every member of  $M_{PQD}(1/2, 1/2; 1/3, 1/3, 1/3)$  is a convex combination of these four extreme points.

Example 3  $m = 3$  and  $n = 4$ .

Let  $p_1, p_2, p_3, q_1, q_2, q_3, q_4$  be seven positive numbers satisfying  $p_1 + p_2 + p_3 = 1 = q_1 + q_2 + q_3 + q_4$ . If  $(p_{ij}) \in M_{PQD}(p_1, p_2, p_3; q_1, q_2, q_3, q_4)$ , then

$$(1) \quad p_3 q_4 \leq p_{34} \leq p_3 \wedge q_4,$$

$$(2) \quad p_{34} \vee (p_3 q_3 + p_3 q_4) \leq p_{33} + p_{34} \leq p_3 \wedge (q_3 + p_{34}),$$

$$(3) \quad p_{34} \vee (p_2 q_4 + p_3 q_4) \leq p_{24} + p_{34} \leq q_4 \wedge (p_2 + p_{34}),$$

$$(4) \quad (p_{33} + p_{34}) \vee (p_3 q_2 + p_3 q_3 + p_3 q_4) \leq p_{32} + p_{33} + p_{34} \leq p_3 \wedge (q_2 + p_{33} + p_{34}),$$

$$(5) \quad (p_{33} + p_{24} + p_{34}) \vee (p_2 q_3 + p_2 q_4 + p_3 q_3 + p_3 q_4) \leq p_{23} + p_{24} + p_{33} + p_{34} \leq (p_2 + p_{33} + p_{34}) \wedge (q_3 + p_{24} + p_{34}),$$

and

$$(6) \quad (p_{23} + p_{24} + p_{34} + p_{32} + p_{33}) \vee (p_2 q_2 + p_2 q_3 + p_2 q_4 + p_3 q_2 + p_3 q_3 + p_3 q_4) \leq p_{22} + p_{23} + p_{24} + p_{32} + p_{33} + p_{34} \leq (p_2 + p_{32} + p_{33} + p_{34}) \wedge (q_2 + p_{23} + p_{24} + p_{33} + p_{34}).$$

Conversely, if  $p_{34}, p_{33}, p_{24}, p_{32}, p_{23}, p_{22}$  are six numbers satisfying the above inequalities, then these six numbers determine uniquely a member of  $M_{PQD}(p_1, p_2, p_3; q_1, q_2, q_3, q_4)$ . We explain how to enumerate all extreme points of the convex set in the simple example  $p_1 = p_2 = p_3 = 1/3$  and  $q_1 = q_2 = q_3 = q_4 = 1/4$ . (The technique in the general case is similar to this special example.) The six inequalities above now become

- (1)  $1/12 \leq p_{34} \leq 3/12,$
- (2)  $p_{34} \vee 2/12 \leq p_{33} + p_{34} \leq 4/12 \wedge (3/12 + p_{34}) = 4/12,$
- (3)  $p_{34} \vee 2/12 \leq p_{24} + p_{34} \leq 3/12 \wedge (4/12 + p_{34}) = 3/12,$
- (4)  $(p_{33} + p_{34}) \vee 3/12 \leq p_{32} + p_{33} + p_{34} \leq 4/12 \wedge (3/12 + p_{33} + p_{34}) = 4/12,$
- (5)  $(p_{33} + p_{24} + p_{34}) \vee 4/12 \leq p_{23} + p_{24} + p_{33} + p_{34} \leq (4/12 + p_{33} + p_{34}) \wedge (3/12 + p_{24} + p_{34}) = 3/12 + p_{24} + p_{34},$
- (6)  $(p_{23} + p_{32} + p_{33} + p_{24} + p_{34}) \vee 6/12 \leq p_{22} + p_{23} + p_{24} + p_{32} + p_{33} + p_{34} \leq (4/12 + p_{32} + p_{33} + p_{34}) \wedge (3/12 + p_{23} + p_{24} + p_{33} + p_{34}).$

The first step in the determination of extreme points is to get rid of the maximum and minimum symbols by splitting some or all inequalities above. For example, inequality (1) can be written as  $1/12 \leq p_{34} \leq 2/12$  and  $2/12 \leq p_{34} \leq 3/12$ . Inequality (3) can be written as  $p_{34} \vee 2/12 \leq p_{33} + p_{34} \leq 3/12$  and  $3/12 \leq p_{33} + p_{34} \leq 4/12$ . The above set of inequalities are equivalent to the following four sets of inequalities.

- I
- (1)  $1/12 \leq p_{34} \leq 2/12$
  - (2)  $2/12 \leq p_{33} + p_{34} \leq 3/12$
  - (3)  $2/12 \leq p_{24} + p_{34} \leq 3/12$
  - (4)  $3/12 \leq p_{32} + p_{33} + p_{34} \leq 4/12$
  - (5) Same as above
  - (6) Same as above

- II (1)  $1/12 \leq p_{34} \leq 2/12$   
(2)  $3/12 \leq p_{33} + p_{34} \leq 4/12$   
(3)  $2/12 \leq p_{24} + p_{34} \leq 3/12$   
(4)  $p_{33} + p_{34} \leq p_{32} + p_{33} + p_{34} \leq 4/12$   
(5) Same as above  
(6) Same as above

- III (1)  $2/12 \leq p_{34} \leq 3/12$   
(2)  $p_{34} \leq p_{33} + p_{34} \leq 3/12$   
(3)  $p_{34} \leq p_{24} + p_{34} \leq 3/12$   
(4)  $3/12 \leq p_{32} + p_{33} + p_{34} \leq 4/12$   
(5) Same as above  
(6) Same as above

- IV (1)  $2/12 \leq p_{34} \leq 3/12$   
(2)  $3/12 \leq p_{33} + p_{34} \leq 4/12$   
(3)  $p_{34} \leq p_{24} + p_{34} \leq 3/12$   
(4)  $p_{33} + p_{34} \leq p_{32} + p_{33} + p_{34} \leq 4/12$   
(5) Same as above  
(6) Same as above

The maximum and minimum symbols in inequalities (5) and (6) stay put in spite of splitting the inequalities (1) and (2). In order to neutralize these symbols, we introduce the following auxiliary inequalities.

$$(2,3) \quad p_{33} + p_{24} + p_{34} \leq 4/12$$

or

$$4/12 \leq p_{33} + p_{24} + p_{34}$$

$$(2,3,4,5) \quad p_{23} + p_{32} + p_{33} + p_{24} + p_{34} \leq 6/12$$

or

$$6/12 \leq p_{23} + p_{32} + p_{33} + p_{24} + p_{34}$$

$$(4,5) \quad 4/12 + p_{32} + p_{33} + p_{34} \leq 3/12 + p_{23} + p_{24} + p_{33} + p_{34}$$

or

$$3/12 + p_{23} + p_{24} + p_{33} + p_{34} \leq 4/12 + p_{32} + p_{33} + p_{34}.$$

Now, a choice of each of the auxiliary inequalities (2,3), (2,3,4,5) and (4,5) is appended to each set of the inequalities I, II, III and IV. This would generate 32 sets of inequalities equivalent to the four sets I, II, III and IV of inequalities. To save space, we will not reproduce these 32 sets of inequalities. A sample set of inequalities is produced below for further discussion.

A sample set of inequalities chosen from 32 sets of inequalities described above.

- (1)  $2/12 \leq p_{34} \leq 3/12$
- (2)  $3/12 \leq p_{33} + p_{34} \leq 4/12$
- (3)  $p_{34} \leq p_{24} + p_{34} \leq 3/12$
- (2,3)  $p_{33} + p_{24} + p_{34} \leq 4/12$
- (4)  $p_{33} + p_{34} \leq p_{32} + p_{33} + p_{34} \leq 4/12$
- (2,3,4,5)  $6/12 \leq p_{23} + p_{32} + p_{33} + p_{24} + p_{34}$
- (5)  $4/12 \leq p_{23} + p_{24} + p_{33} + p_{34} \leq 3/12 + p_{24} + p_{34}$
- (4,5)  $4/12 + p_{32} + p_{33} + p_{34} \leq 3/12 + p_{23} + p_{24} + p_{33} + p_{34}$
- (6)  $p_{23} + p_{32} + p_{33} + p_{24} + p_{34} \leq p_{22} + p_{23} + p_{24} + p_{32} + p_{33} + p_{34}$   
 $\leq 4/12 + p_{32} + p_{33} + p_{34}.$

The above set of inequalities is obtained from the set IV of inequalities by appending the first choice of (2,3), the second choice of (2,3,4,5) and the first choice of (4,5).

In order to obtain a member of  $M_{p_{QD}}(1/3,1/3,1/3; 1/4,1/4,1/4,1/4)$  we proceed as follows. Set the central expression in each of the main six inequalities equal to the quantity either on the left or the right of the inequalities and then solve the system of equations thus arise in the unknowns  $p_{34}, p_{33}, p_{24}, p_{32}, p_{23}, p_{22}$  making sure that the constraints imposed by the auxiliary inequalities are satisfied. These system of equations are easy to solve. The solution will give a member of  $M_{p_{QD}}(1/3,1/3,1/3; 1/4,1/4,1/4,1/4)$ .

Generate members of  $M_{PQD}(1/3,1/3,1/3; 1/4,1/4,1/4,1/4)$  by following the above procedure for each set of the 32 sets of inequalities. The set of extreme points of the convex set of interest is a subset of these solutions. There will be a large amount of duplicates and some of the solutions obtained are already convex combinations of other solutions. After considerable amount of weeding, we got the following matrices as the entire collection of extreme points of the convex set  $M_{PQD}(1/3,1/3,1/3; 1/4,1/4,1/4,1/4)$ .

$$1. \begin{bmatrix} 1/12 & 1/12 & 1/12 & 1/12 \\ 1/12 & 1/12 & 1/12 & 1/12 \\ 1/12 & 1/12 & 1/12 & 1/12 \end{bmatrix}$$

$$2. \begin{bmatrix} 2/12 & 0 & 1/12 & 1/12 \\ 0 & 2/12 & 1/12 & 1/12 \\ 1/12 & 1/12 & 1/12 & 1/12 \end{bmatrix}$$

$$3. \begin{bmatrix} 1/12 & 1/12 & 1/12 & 1/12 \\ 2/12 & 0 & 1/12 & 1/12 \\ 0 & 2/12 & 1/12 & 1/12 \end{bmatrix}$$

$$4. \begin{bmatrix} 2/12 & 0 & 1/12 & 1/12 \\ 1/12 & 1/12 & 1/12 & 1/12 \\ 0 & 2/12 & 1/12 & 1/12 \end{bmatrix}$$

$$5. \begin{bmatrix} 3/12 & 0 & 0 & 1/12 \\ 0 & 1/12 & 2/12 & 1/12 \\ 0 & 2/12 & 1/12 & 1/12 \end{bmatrix}$$

$$6. \begin{bmatrix} 1/12 & 2/12 & 0 & 1/12 \\ 1/12 & 0 & 2/12 & 1/12 \\ 1/12 & 1/12 & 1/12 & 1/12 \end{bmatrix}$$

$$7. \begin{bmatrix} 2/12 & 1/12 & 0 & 1/12 \\ 0 & 1/12 & 2/12 & 1/12 \\ 1/12 & 1/12 & 1/12 & 1/12 \end{bmatrix}$$

$$8. \begin{bmatrix} 3/12 & 1/12 & 0 & 0 \\ 0 & 0 & 2/12 & 2/12 \\ 0 & 2/12 & 1/12 & 1/12 \end{bmatrix}$$

$$9. \begin{bmatrix} 2/12 & 0 & 2/12 & 0 \\ 1/12 & 1/12 & 0 & 2/12 \\ 0 & 2/12 & 1/12 & 1/12 \end{bmatrix}$$

$$10. \begin{bmatrix} 1/12 & 1/12 & 2/12 & 0 \\ 2/12 & 0 & 0 & 2/12 \\ 0 & 2/12 & 1/12 & 1/12 \end{bmatrix}$$



$$11. \begin{bmatrix} 1/12 & 2/12 & 0 & 1/12 \\ 2/12 & 1/12 & 0 & 1/12 \\ 0 & 0 & 3/12 & 1/12 \end{bmatrix}$$

$$12. \begin{bmatrix} 3/12 & 0 & 0 & 1/12 \\ 0 & 3/12 & 0 & 1/12 \\ 0 & 0 & 3/12 & 1/12 \end{bmatrix}$$

$$13. \begin{bmatrix} 1/12 & 0 & 2/12 & 1/12 \\ 1/12 & 2/12 & 1/12 & 0 \\ 1/12 & 1/12 & 0 & 2/12 \end{bmatrix}$$

$$14. \begin{bmatrix} 1/12 & 2/12 & 0 & 1/12 \\ 1/12 & 0 & 3/12 & 0 \\ 1/12 & 1/12 & 0 & 2/12 \end{bmatrix}$$

$$15. \begin{bmatrix} 2/12 & 1/12 & 0 & 1/12 \\ 0 & 1/12 & 3/12 & 0 \\ 1/12 & 1/12 & 0 & 2/12 \end{bmatrix}$$

$$16. \begin{bmatrix} 1/12 & 1/12 & 1/12 & 1/12 \\ 2/12 & 0 & 2/12 & 0 \\ 0 & 2/12 & 0 & 2/12 \end{bmatrix}$$

$$17. \begin{bmatrix} 2/12 & 0 & 1/12 & 1/12 \\ 1/12 & 1/12 & 2/12 & 0 \\ 0 & 2/12 & 0 & 2/12 \end{bmatrix}$$

$$18. \begin{bmatrix} 2/12 & 1/12 & 0 & 1/12 \\ 1/12 & 0 & 3/12 & 0 \\ 0 & 2/12 & 0 & 2/12 \end{bmatrix}$$

$$19. \begin{bmatrix} 1/12 & 2/12 & 1/12 & 0 \\ 1/12 & 0 & 2/12 & 1/12 \\ 1/12 & 1/12 & 0 & 2/12 \end{bmatrix}$$

$$20. \begin{bmatrix} 2/12 & 1/12 & 1/12 & 0 \\ 0 & 1/12 & 2/12 & 1/12 \\ 1/12 & 1/12 & 0 & 2/12 \end{bmatrix}$$

$$21. \begin{bmatrix} 2/12 & 2/12 & 0 & 0 \\ 0 & 0 & 3/12 & 1/12 \\ 1/12 & 1/12 & 0 & 2/12 \end{bmatrix}$$

$$22. \begin{bmatrix} 2/12 & 0 & 2/12 & 0 \\ 1/12 & 1/12 & 1/12 & 1/12 \\ 0 & 2/12 & 0 & 2/12 \end{bmatrix}$$

$$23. \begin{bmatrix} 1/12 & 1/12 & 2/12 & 0 \\ 2/12 & 0 & 1/12 & 1/12 \\ 0 & 2/12 & 0 & 2/12 \end{bmatrix}$$

$$24. \begin{bmatrix} 1/12 & 1/12 & 1/12 & 1/12 \\ 1/12 & 1/12 & 2/12 & 0 \\ 1/12 & 1/12 & 0 & 2/12 \end{bmatrix}$$

$$25. \begin{bmatrix} 2/12 & 0 & 1/12 & 1/12 \\ 0 & 2/12 & 2/12 & 0 \\ 1/12 & 1/12 & 0 & 2/12 \end{bmatrix}$$

$$26. \begin{bmatrix} 2/12 & 0 & 1/12 & 1/12 \\ 1/12 & 1/12 & 2/12 & 0 \\ 0 & 2/12 & 0 & 2/12 \end{bmatrix}$$

$$27. \begin{bmatrix} 1/12 & 1/12 & 2/12 & 0 \\ 0 & 2/12 & 1/12 & 1/12 \\ 2/12 & 0 & 0 & 2/12 \end{bmatrix}$$

$$28. \begin{bmatrix} 1/12 & 3/12 & 0 & 0 \\ 0 & 0 & 3/12 & 1/12 \\ 2/12 & 0 & 0 & 2/12 \end{bmatrix}$$

$$29. \begin{bmatrix} 2/12 & 0 & 1/12 & 1/12 \\ 1/12 & 3/12 & 0 & 0 \\ 0 & 0 & 2/12 & 2/12 \end{bmatrix}$$

$$30. \begin{bmatrix} 1/12 & 2/12 & 0 & 1/12 \\ 2/12 & 1/12 & 1/12 & 0 \\ 0 & 0 & 2/12 & 2/12 \end{bmatrix}$$

$$31. \begin{bmatrix} 0 & 2/12 & 1/12 & 1/12 \\ 3/12 & 1/12 & 0 & 0 \\ 0 & 0 & 2/12 & 2/12 \end{bmatrix}$$

$$32. \begin{bmatrix} 1/12 & 1/12 & 2/12 & 0 \\ 1/12 & 2/12 & 1/12 & 0 \\ 1/12 & 0 & 0 & 3/12 \end{bmatrix}$$

$$33. \begin{bmatrix} 2/12 & 0 & 2/12 & 0 \\ 0 & 3/12 & 1/12 & 0 \\ 1/12 & 0 & 0 & 3/12 \end{bmatrix}$$

$$34. \begin{bmatrix} 1/12 & 3/12 & 0 & 0 \\ 1/12 & 0 & 3/12 & 0 \\ 1/12 & 0 & 0 & 3/12 \end{bmatrix}$$

$$35. \begin{bmatrix} 1/12 & 0 & 3/12 & 0 \\ 2/12 & 2/12 & 0 & 0 \\ 0 & 1/12 & 0 & 3/12 \end{bmatrix}$$

$$36. \begin{bmatrix} 1/12 & 2/12 & 1/12 & 0 \\ 2/12 & 0 & 2/12 & 0 \\ 0 & 1/12 & 0 & 3/12 \end{bmatrix}$$

$$37. \begin{bmatrix} 3/12 & 0 & 1/12 & 0 \\ 0 & 2/12 & 2/12 & 0 \\ 0 & 1/12 & 0 & 3/12 \end{bmatrix}$$

$$38. \begin{bmatrix} 1/12 & 3/12 & 0 & 0 \\ 2/12 & 0 & 2/12 & 0 \\ 0 & 0 & 1/12 & 3/12 \end{bmatrix}$$

$$39. \begin{bmatrix} 2/12 & 2/12 & 0 & 0 \\ 1/12 & 1/12 & 2/12 & 0 \\ 0 & 0 & 1/12 & 3/12 \end{bmatrix}.$$

#### 4. Testing independence against strict positive quadrant dependence

Let  $X$  and  $Y$  be two random variables with known marginal distributions and unknown joint distribution. We want to test the hypothesis that  $X$  and  $Y$  are independent against the hypothesis that they are strictly positive quadrant dependent. By strict positive quadrant dependence we mean positive quadrant dependence but not independence. The data consist of  $N$  independent realizations of the vector  $(X,Y)$ . Let  $\tau$  be a test proposed for testing the hypothesis of independence based on the given data. Let  $\lambda$  be the distribution of  $X$  and  $\nu$  that of  $Y$ . Let  $M_{PQD}(\lambda,\nu)$  be the collection of all bivariate distributions with fixed marginals  $\lambda$  and  $\nu$  which are positive quadrant dependent. The power function of the test  $\tau$  can be defined formally as follows.

$$B_{\tau}(\mu) = \Pr\{\tau \text{ rejects the null hypothesis} / \mu\}$$

for  $\mu$  in  $M_{PQD}(\lambda,\nu)$ . The above probability is computed when the joint

distribution of  $X$  and  $Y$  is  $\mu$ . The calculation of the power function of the test  $\tau$  whose domain of definition is  $M_{PQD}(\lambda, \nu)$  is very tedious. Moreover, if we wish to compare the performance of two tests to discriminate the null hypothesis of independence against the alternative hypothesis of strict positive quadrant dependence, we need to compare their power functions. This comparison then becomes doubly more difficult to achieve. But the following theorem asserts that it suffices to compare the powers at the extreme points of  $M_{PQD}(\lambda, \nu)$  only.

Theorem 5 Let  $\mu, \mu^1, \mu^2, \dots, \mu^k$  be members of  $M_{PQD}(\lambda, \nu)$  such that  $\mu = \alpha_1 \mu^1 + \alpha_2 \mu^2 + \dots + \alpha_k \mu^k$  for some  $\alpha_1, \alpha_2, \dots, \alpha_k \geq 0$  with  $\sum_{i=1}^k \alpha_i = 1$ . Then

$$\beta_{\tau}(\mu) = \sum_{i=1}^k \alpha_i \beta_{\tau}(\mu^i)$$

The above result can be used as follows. Suppose each of  $X$  and  $Y$  takes finitely many values. Then the convex set  $M_{PQD}(\lambda, \nu)$  has finitely many extreme points and every member of  $M_{PQD}(\lambda, \nu)$  can be written as a convex combination of these extreme points. Then the power of the test  $\tau$

evaluated at any given  $\mu$  in  $M_{PQD}(\lambda, \nu)$  is precisely the same convex combination of the powers of the test evaluated at each of the extreme points. This result also points out that in order to compare the performance of two given tests, it suffices to compare the powers evaluated at the extreme points. As an illustration, we consider the case when  $X$  takes values 1 and 2, and  $Y$  takes values 1, 2 and 3. Let  $n_{ij}$  = total number of  $(X, Y)$ 's with  $X = i$  and  $Y = j$ ,  $i = 1, 2$  and  $j = 1, 2, 3$ . The data can be arranged in the form of a contingency table as follows.

$$\begin{bmatrix} n_{11} & n_{12} & n_{13} \\ n_{21} & n_{22} & n_{23} \end{bmatrix}$$

In this section, we compare the performance of two tests for testing the null hypothesis of independence against the alternative of strict positive quadrant dependence in the context of  $2 \times 3$  contingency tables above.

#### $T_1$ : Test based-on gamma ratio

Let the bivariate distribution of  $X$  and  $Y$  be given by

$$\begin{bmatrix} p_{11} & p_{12} & p_{13} \\ p_{21} & p_{22} & p_{23} \end{bmatrix} .$$

The Gamma Ratio (see Goodman and Kruskal [25]) of  $X$  and  $Y$  is defined by

$$\gamma = \frac{\pi_c - \pi_d}{\pi_c + \pi_d} ,$$

where  $\pi_c = 2p_{11}(p_{22} + p_{23}) + 2p_{12}p_{13}$  and  $\pi_d = 2p_{13}(p_{21} + p_{22}) + 2p_{12}p_{21}$ .

One can show that  $\gamma = 0$  if  $X$  and  $Y$  are independent. See

Agresti (1984, p.160). One can also show that  $\gamma \geq 0$  if  $X$  and  $Y$  are positive quadrant dependent. An estimate of  $\gamma$  based on the sample given above is given by

$$\hat{\gamma} = \frac{C - D}{C + D},$$

where  $C$  = The total number of concordant pairs =  $n_{11}(n_{22} + n_{23}) + n_{12}n_{23}$   
and  $D$  = The total number of discordant pairs =  $n_{13}(n_{21} + n_{22}) + n_{12}n_{21}$ .

The following is a natural test based on  $\hat{\gamma}$  for testing the above null hypothesis against the specific alternative mentioned thereby.

Test  $T_1$  : Reject the null hypothesis if  $\hat{\gamma} \geq a$ .

$T_2$  : Test based on eigen values

Let the marginal distribution of  $X$  be given by  $p_1$  and  $p_2$ , and that of  $Y$  by  $q_1, q_2$  and  $q_3$ . Let

$$Q = \begin{bmatrix} \frac{p_{11}}{\sqrt{p_1 q_1}} & \frac{p_{12}}{\sqrt{p_1 q_2}} & \frac{p_{13}}{\sqrt{p_1 q_3}} \\ \frac{p_{21}}{\sqrt{p_2 q_1}} & \frac{p_{22}}{\sqrt{p_2 q_2}} & \frac{p_{23}}{\sqrt{p_2 q_3}} \end{bmatrix}.$$

Let  $\kappa_1$  and  $\kappa_2$  be the eigen values of  $QQ^T$ , where  $T$  denotes operation transpose on matrices. We give some properties of these eigen values below. For further details, see Lancaster [11] and [12], O'Neill ([17], [18], and [19]).

### Properties

1. One of the eigen values is always equals unity. Let us use  $\kappa_1$  for this eigen value.
2. If  $X$  and  $Y$  are independent,  $\kappa_2 = 0$ .
3. If  $X$  and  $Y$  are strictly positive quadrant dependent, then  $\kappa_2 > 0$ .

We estimate  $\kappa_1 + \kappa_2$  based on the data given above as follows.

Let

$$B = \begin{bmatrix} \frac{n_{11}}{N\sqrt{p_1q_1}} & \frac{n_{12}}{N\sqrt{p_1q_2}} & \frac{n_{13}}{N\sqrt{p_1q_3}} \\ \frac{n_{21}}{N\sqrt{p_2q_1}} & \frac{n_{22}}{N\sqrt{p_2q_2}} & \frac{n_{23}}{N\sqrt{p_2q_3}} \end{bmatrix}.$$

Let  $\hat{\kappa}_1^2$  and  $\hat{\kappa}_2^2$  be the eigen values of  $BB^T$ . Then we propose  $\hat{\kappa} = \hat{\kappa}_1 + \hat{\kappa}_2$  as an estimator of  $\kappa_1 + \kappa_2$ .

Test  $T_2$  : Reject the null hypothesis if  $\hat{\kappa} \geq a$ .

We discuss the performance of these two tests in the case of two specific examples given below.

Example 1  $p_1 = p_2 = 1/2$  and  $q_1 = q_2 = q_3 = 1/3$ .

Example 2  $p_1 = 1/4, p_2 = 3/4$  and  $q_1 = 1/2, q_2 = 1/4, q_3 = 1/4$ .

Let us now elaborate on some of the properties of the eigen values of  $QQ^T$ . The second eigen value of  $QQ^T$ ,  $\kappa_2$ , can be worked out explicitly.

$$\kappa_2 = \frac{(p_{11}p_{22} - p_{12}p_{21})^2}{p_1p_2q_1q_2} + \frac{(p_{11}p_{23} - p_{13}p_{21})^2}{p_1p_2q_1q_3} + \frac{(p_{12}p_{23} - p_{13}p_{22})^2}{p_1p_2q_2q_3}.$$

From this it follows that  $\kappa_2 = 0$  if and only if  $X$  and  $Y$  are independent.

At this juncture, we want to make some remarks on the definition of the matrix  $B$  above. In order to develop an estimator of  $\kappa_2$ , it is natural to divide each frequency  $n_{ij}$  in  $B$  by the square root of the product of the corresponding marginal totals  $n_{i.}$  and  $n_{.j}$ . See O'Neill ([17], [18], [19]). If we had proceeded as outlined by O'Neill, one of the eigen values of  $BB^T$  would always be equal to unity. In our definition of  $B$ , it is not true that one of the eigen values of  $BB^T$  would always be equal to unity. Since we know the marginal probability laws of  $X$  and  $Y$ , we need to estimate only  $p_{ij}$ 's by  $n_{ij}/N$ 's. This what motivated us to define the matrix  $B$  the way it was defined above. However, in view of the above formula for  $\kappa_2$ , one could estimate  $\kappa_2$  directly without having to define the matrix  $B$ . Accordingly, let

$$\tilde{\kappa}_2 = \frac{(n_{11}n_{22} - n_{12}n_{21})^2}{N^4 p_1p_2q_1q_2} + \frac{(n_{11}n_{23} - n_{12}n_{21})^2}{N^4 p_1p_2q_1q_3} + \frac{(n_{12}n_{23} - n_{13}n_{22})^2}{N^4 p_1p_2q_2q_3}.$$



We can build a test based on the statistic  $\tilde{\kappa}_2$ .

Test  $T_3$ : Reject the null hypothesis if  $\tilde{\kappa}_2 \geq a$ .

We discuss the performance of these three tests in the case of two specific examples given below.

Example 1  $p_1 = p_2 = 1/2$  and  $q_1 = q_2 = q_3 = 1/3$ .

Example 2  $p_1 = 1/4$ ,  $p_2 = 3/4$  and  $q_1 = 1/2$ ,  $q_2 = 1/4$ ,  $q_3 = 1/4$ .

The performance of  $T_1$  and  $T_3$  was compared in detail in Subramanyam and Bhaskara Rao [12]. We now compare the performance of the tests  $T_1$ ,  $T_2$  and  $T_3$  together under the level of significance  $\alpha = 0.01, 0.025, 0.05$  and  $N = 15, 20$  and  $25$ . The exact distribution of  $\hat{\gamma}$ ,  $\hat{\kappa}$  and  $\hat{\kappa}_2$  is evaluated for each of the sample sizes  $N = 15, 20$  and  $25$  and the power of the tests  $T_1$ ,  $T_2$  and  $T_3$  is evaluated at each of the extreme point distributions of the above examples using these exact distributions. The graphs\* of these distributions are given at the end of this section. Nguyen and Sampson [8] evaluated the powers of tests based on some other statistics at some specific alternative distributions by simulating these distributions.

\*The authors wish to thank Ron Chao for his valuable help in the computations.

Example No.	Extreme Points	$\gamma$	$\kappa_2$
1.	$P_1 = \begin{bmatrix} 1/6 & 1/6 & 1/6 \\ 1/6 & 1/6 & 1/6 \end{bmatrix}$	0	0
	$P_2 = \begin{bmatrix} 1/6 & 1/3 & 0 \\ 1/6 & 0 & 1/3 \end{bmatrix}$	1/2	2/3
	$P_3 = \begin{bmatrix} 1/3 & 1/6 & 0 \\ 0 & 1/6 & 1/3 \end{bmatrix}$	1	2/3
	$P_4 = \begin{bmatrix} 1/3 & 0 & 1/6 \\ 0 & 1/3 & 1/6 \end{bmatrix}$	1/2	2/3
2.	$P_5 = \begin{bmatrix} 2/16 & 1/16 & 1/16 \\ 6/16 & 3/16 & 3/16 \end{bmatrix}$	0	0
	$P_6 = \begin{bmatrix} 2/16 & 2/16 & 0 \\ 6/16 & 2/16 & 4/16 \end{bmatrix}$	1/4	1/6
	$P_7 = \begin{bmatrix} 4/16 & 0 & 0 \\ 4/16 & 4/16 & 4/16 \end{bmatrix}$	1	1/3
	$P_8 = \begin{bmatrix} 3/16 & 0 & 1/16 \\ 5/16 & 4/16 & 3/16 \end{bmatrix}$	4/10	1/8

Table 1: Power functions of the tests  $T_1$ ,  $T_2$ , and  $T_3$ .

Sample size: N	15			20			25		
Distribution	$\beta_{T_1}(\cdot)$	$\beta_{T_2}(\cdot)$	$\beta_{T_3}(\cdot)$	$\beta_{T_1}(\cdot)$	$\beta_{T_2}(\cdot)$	$\beta_{T_3}(\cdot)$	$\beta_{T_1}(\cdot)$	$\beta_{T_2}(\cdot)$	$\beta_{T_3}(\cdot)$
Example 1									
$P_1$ (size)	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
$P_2$	0.0768	0.3944	0.5021	0.1389	0.6477	0.6892	0.2080	0.8817	0.8339
$P_3$	0.9999	0.3944	0.5360	1.0000	0.6477	0.8276	1.0000	0.8817	0.9569
$P_4$	0.0768	0.3944	0.5360	0.1389	0.6477	0.8276	0.2080	0.8817	0.9569
Critical value: $\alpha$	0.90	1.99	0.66	0.78	1.73	0.49	0.71	1.60	0.39
Example 2									
$P_5$ (size)	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
$P_6$	0.1224	0.0540	0.1037	0.0661	0.0821	0.1360	0.0348	0.1144	0.1670
$P_7$	0.9866	0.1110	0.0923	0.9968	0.1824	0.1855	0.9992	0.2501	0.3131
$P_8$	0.3664	0.0403	0.0194	0.2719	0.0548	0.0348	0.1984	0.0654	0.0595
Critical value: $\alpha$	1.00	2.05	0.77	1.00	1.78	0.56	1.00	1.62	0.44

Table 2: Power functions of the tests  $T_1$ ,  $T_2$ , and  $T_3$ .

Sample size: N	15			20			25		
Distribution	$\beta_{T_1}(\cdot)$	$\beta_{T_2}(\cdot)$	$\beta_{T_3}(\cdot)$	$\beta_{T_1}(\cdot)$	$\beta_{T_2}(\cdot)$	$\beta_{T_3}(\cdot)$	$\beta_{T_1}(\cdot)$	$\beta_{T_2}(\cdot)$	$\beta_{T_3}(\cdot)$
Example 1									
$P_1$ (size)	0.025	0.025	0.025	0.025	0.025	0.025	0.025	0.025	0.025
$P_2$	0.1733	0.6165	0.6581	0.2602	0.8693	0.8241	0.3334	1.0000	0.9365
$P_3$	0.9999	0.6165	0.7817	1.0000	0.8693	0.9467	1.0000	1.0000	0.9925
$P_4$	0.1733	0.6165	0.7817	0.2602	0.8693	0.9467	0.3334	1.0000	0.9925
Critical value: a	0.80	1.83	0.51	0.69	1.61	0.38	0.62	1.50	0.30
Example 2									
$P_5$ (size)	0.025	0.025	0.025	0.025	0.025	0.025	0.025	0.025	0.025
$P_6$	0.1224	0.1181	0.1747	0.0661	0.1492	0.2139	0.0348	0.1884	0.2569
$P_7$	0.9866	0.2152	0.1993	0.9968	0.2802	0.3508	0.9993	0.4071	0.5017
$P_8$	0.3664	0.0965	0.0571	0.2719	0.1047	0.3508	0.1984	0.1352	0.1500
Critical value: a	1.00	1.87	0.56	1.00	1.66	0.42	1.00	1.52	0.33









Table 3: Power functions of the tests  $T_1$ ,  $T_2$  and  $T_3$

Sample size: N	15			20			25		
Distribution	$\beta_{T_1}()$	$\beta_{T_2}()$	$\beta_{T_3}()$	$\beta_{T_1}()$	$\beta_{T_2}()$	$\beta_{T_3}()$	$\beta_{T_1}()$	$\beta_{T_2}()$	$\beta_{T_3}()$
$P_1$ (size)	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05
$P_2$	0.3229	0.7602	0.7771	0.3927	0.9964	0.9209	0.4761	0.9999	0.9768
$P_3$	0.9999	0.7602	0.8927	1.0000	0.9964	0.9822	1.0000	0.9999	0.9976
$P_4$	0.2936	0.7602	0.8927	0.4142	0.9964	0.9822	0.4761	0.9999	0.9976
Critical value: a	0.69	1.73	0.40	0.60	1.52	0.30	0.53	1.43	0.24
$P_5$ (size)	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05
$P_6$	0.1224	0.1870	0.2511	0.0661	0.2312	0.3144	0.0780	0.3030	0.3524
$P_7$	0.9866	0.3286	0.3526	0.9968	0.4447	0.5357	0.9992	0.5727	0.6650
$P_8$	0.3667	0.1615	0.1279	0.2719	0.1969	0.2143	0.2399	0.2366	0.2847
Critical value: a	1.00	1.74	0.43	1.00	1.55	0.31	0.73	1.44	0.25









### Conclusions

1. The power of the test  $T_1$  dominates the power of the other two tests at the extreme point distribution  $P_3$  in Example 1 and  $P_7$  in Example 2. This is not surprising as the Gamma Ratio achieves the perfect value unity under  $P_3$  in Example 1 and  $P_7$  in Example 2.
2. On the whole  $T_2$  seems to perform well in comparison with the other two tests. Even under the distribution  $P_3$  in Example 1 and  $P_7$  in Example 2,  $T_2$  is not overpowered by  $T_1$ .
3. Some extensive studies are needed to be carried out to see whether  $T_2$  is preferable to the other two under different sets of marginal distributions.

GRAPHS OF THE DISTRIBUTIONS OF  $\hat{\gamma}$  FOR  $N=10,15,20,25$  UNDER  $P_1$  AND  $P_2$ :











GRAPHS OF THE DISTRIBUTIONS OF  $\hat{\gamma}$  FOR  $N=10,15,20,25$  UNDER  $P_3$  AND  $P_4$ .





















GRAPHS OF THE DISTRIBUTIONS OF  $\hat{\gamma}$  FOR  $N=10,15,20,25$  UNDER  $P_5$  AND  $P_6$ .











GRAPHS OF THE DISTRIBUTIONS OF  $\hat{\gamma}$  FOR  $N=10,15,20,25$  UNDER  $P_7$  AND  $P_8$ .







GRAPHS OF THE DISTRIBUTIONS OF  $\hat{\kappa} = \hat{\kappa}_1 + \hat{\kappa}_2$  FOR  $N = 10, 15, 20, 25$  UNDER  $P_1$  AND  $P_2$ .











GRAPHS OF THE DISTRIBUTIONS OF  $\hat{\kappa} = \hat{\kappa}_1 + \hat{\kappa}_2$  FOR  $N = 10, 15, 20, 25$  UNDER  $P_3$  AND  $P_4$ .





















GRAPHS OF THE DISTRIBUTIONS OF  $\hat{\kappa} = \hat{\kappa}_1 + \hat{\kappa}_2$  FOR  $N = 10, 15, 20, 25$  UNDER  $P_7$  and  $P_8$ .



GRAPHS OF THE DISTRIBUTIONS OF  $\hat{\kappa}_2$  FOR  $N = 10, 15, 20, 25$  UNDER  $P_1$  AND  $P_2$ .











GRAPHS OF THE DISTRIBUTIONS OF  $\hat{\kappa}_2$  FOR  $N = 10, 15, 20, 25$  UNDER  $P_3$  AND  $P_4$ .













GRAPHS OF THE DISTRIBUTIONS OF  $\hat{\kappa}_2$  FOR  $N = 10, 15, 20, 25$  UNDER  $P_5$  and  $P_6$ .



GRAPHS OF THE DISTRIBUTIONS OF  $\hat{\kappa}_2$  FOR  $N = 10, 15, 20, 25$  UNDER  $P_7$  and  $P_8$ .



### 5. Inference on the structure of dependence

In two-way contingency tables, the  $\chi^2$  test for independence has been widely used. When the test for independence is rejected, it is of interest to study the structure of dependence between the  $a+1$  rows and  $b+1$  columns. In this section, we write the matrices  $F = (f_{ij})$  and  $B = (b_{ij})$  in terms of their eigenvalues and eigenvectors by singular value decomposition; here  $f_{ij} = p_{ij}/\sqrt{p_i q_j}$ ,  $b_{ij} = n_{ij}/\sqrt{n_i n_j}$ ,  $p_i = p_{i1} + \dots + p_{i,b+1}$ ,  $q_j = p_{1j} + \dots + p_{a+1,j}$ ,  $n_i = n_{i1} + \dots + n_{i,b+1}$  and  $n_j = n_{1j} + \dots + n_{a+1,j}$ . Taking advantage of the above decomposition, we propose procedures to find out as to which of the last  $t$  eigenvalues of  $FF'$  are zero. The distribution theory associated with the above procedures are also investigated. Some aspects of the above problem were discussed by O'Neill ([17], [18], [19]). The problem of determination of the rank of  $F$  is discussed in a forthcoming paper of Z.D. Bai, P.R. Krishnaiah and L.C. Zhao from the point of view of model selection using an information theoretic criterion. The above authors also established the strong consistency of their procedure. In this section, we use the notation  $n = \sum_{i=1}^{a+1} \sum_{j=1}^{b+1} n_{ij}$  is fixed and the marginal totals  $n_i$  and  $n_j$  are random.

Consider the model

$$p_{ij} = p_i q_j \zeta_{ij} \quad (5.1)$$

$i = 1, 2, \dots, a+1$ ,  $j = 1, 2, \dots, b+1$ . Without losing generality we assume that  $a \leq b$ . Under the above model, it is of interest to test for the structure of  $\zeta_{ij}$ . From singular value decomposition of a matrix, it is known (e.g., see Lancaster [12]) that

$$F = \begin{pmatrix} \xi^* & n^* \\ 0 & 0 \end{pmatrix} \delta_0 + \sum_{u=1}^a \delta_u \begin{pmatrix} \xi_u^* & n_u^* \\ 0 & 0 \end{pmatrix} \quad (5.2)$$

where  $\delta_0 \geq \delta_1 \geq \dots \geq \delta_a$  are the eigenvalues of  $F$ ,  $\delta_0 = 1$ ,  $\xi_u^*$  is the eigenvector of  $FF'$  corresponding to  $\delta_u^2$  and  $\eta_u^*$  is the eigenvector of  $F'F$  corresponding to  $\delta_u^2$ . In (5.2)

$$\begin{aligned}\xi_u^* &= (\sqrt{p_1} \xi_{u1}, \dots, \sqrt{p_{a+1}} \xi_{u,a+1})' \\ \eta_u^* &= (\sqrt{q_1} \eta_{u1}, \dots, \sqrt{q_{b+1}} \eta_{u,b+1})' \\ \xi_{01} &= \dots = \xi_{0,a+1} = 1, \eta_{01} = \dots = \eta_{0,b+1} = 1.\end{aligned}\tag{5.3}$$

We are interested in finding out as to how many  $\delta_u$ 's are equal to zero. This problem is analogous to the problem of studying the structure of interaction term in two-way classification model with one observation per cell. So, we will briefly discuss the above model.

Let

$$E(x_{ij}) = \mu + \alpha_i + \beta_j + \Delta_{ij}\tag{5.4}$$

for  $i = 1, 2, \dots, (a + 1)$ ,  $j = 1, 2, \dots, (b + 1)$ .

$$\sum_i \alpha_i = \sum_j \beta_j = \sum_i \Delta_{ij} = \sum_j \Delta_{ij} = 0.$$

Also  $\mu$ ,  $\alpha_i$ ,  $\beta_j$  and  $\Delta_{ij}$  respectively denote the general mean,  $i$ -th row effect,  $j$ -th column effect, and interaction in  $i$ -th row and  $j$ -th column. Also, let  $\Delta = (\Delta_{ij})$  and  $a < b$ . We assume that  $x_{ij}$ 's are distributed independently and with variance  $\sigma^2$ . The problem of testing the hypothesis  $\Delta = 0$  was first considered by Fisher and MacKenzie [4], and later by Williams [23], Tukey [22] and others when the underlying distribution is normal. Fisher and MacKenzie [4] considered this problem using eigenvalues of certain matrix. For a review of the literature, the reader is

referred to Krishnaiah and Yochmowitz [10]. Now, let  $E(x_{ij}) = \log p_{ij}$ ,  $\mu = \log c$ ,  $\alpha_i = \log p_i$ ,  $\beta_j = \log p_j$  and  $\Delta_{ij} = \log \delta_{ij}$ . Then the model (5.4) can be written as

$$p_{ij} = cp_i q_j \zeta_{ij}. \quad (5.5)$$

But, here we do not assume that the conditions (5.5) are satisfied. But,

$$\log \left( \sum_i p_i \right) = \log \left( \sum_j q_j \right) = 0. \quad (5.6)$$

We may assume that  $c = 1$  (i.e.,  $\mu = 0$ ). We can write (5.6) as

$$p_{ij} = cp_i q_j \exp(\eta_{ij}) \quad (5.7)$$

and express  $\eta = (\eta_{ij})$  in terms of its eigenvalues and eigenvectors using spectral decomposition of a matrix. Then, we can draw inference on the rank of  $\eta$ . This problem is different from the problem of drawing inference on the rank of  $\zeta$  except for the special case when the rank of  $\eta$  is zero. This special case is equivalent to the statement that the rank of  $\zeta$  is one. In studying the interaction term in two-way classification model, Tukey [22] and Mandel [15] assumed certain structures on interaction term. We can assume similar structures on the models (5.6) and (5.7). As far as the models (5.6) and (5.7) are concerned, they are analogous to the well known two-way classification model with interaction and one observation per cell. But, the problems of estimation and distributions of test statistics are of different nature. In general we may also consider models of the form

$$p_{ij} = f(\alpha_i, \beta_j, \eta_{ij})$$

where  $f(\cdot)$  is a suitably chosen function of  $\alpha_i$ ,  $\beta_j$ , and  $\gamma_{ij}$ . For example  $f(x,y,z) = f_1(x)f_2(y)f_3(z)$ . As another possibility, we may also consider the model

$$p_{ij} = \mu + \alpha_i + \beta_j + \gamma_{ij} \quad (5.8)$$

as in two-way classification with interaction.

Goodman [6] discussed the model (5.7) when  $n$  is written in terms of its eigenvalues and eigenvectors. O'Neil ([17], [18], [19]) discussed some aspects of the asymptotic distribution theory associated with finding the rank of the matrix  $\zeta$ . In this section, we propose various test procedures for determination of the rank of  $\zeta$  and investigate some problems on the asymptotic distributions of the test statistics.

We now discuss the problem of testing for the rank of the matrix  $\zeta$ . If we know in advance the rank of  $\zeta$ , we can use that knowledge in estimating the unknown parameters more accurately. For example, the maximum likelihood estimates of  $p_{ij}$ 's when the rank of  $F$  is one are not the same as when the rank of  $F$  is greater than one.

Now, let  $B = (b_{ij})$  where

$$b_{ij} = n_{ij} / \sqrt{n_{i\cdot} n_{\cdot j}} \quad (5.9)$$

Then, using the spectral decomposition of  $B$ , we have

$$B = \hat{\xi}_{\cdot 0}^* \hat{\eta}_{\cdot 0}^{*'} \hat{\delta}_0 + \sum_{u=1}^a \hat{\delta}_u \hat{\xi}_u^* \hat{\eta}_u^{*'} \quad (5.10)$$

where  $\hat{\delta}_0 = 1$ ,

$$\hat{\xi}_u^* = ((n_{1\cdot}/n)^{1/2} \hat{\xi}_{u1}, \dots, (n_{a+1\cdot}/n)^{1/2} \hat{\xi}_{u,a+1}),$$

$$\hat{\eta}_u^* = ((n_{.1}/n)^{1/2} \hat{\eta}_{u1}, \dots, (n_{.b+1}/n)^{1/2} \hat{\eta}_{u,b+1})'$$

$$\hat{\xi}_{01} = \dots = \hat{\xi}_{0,a+1} = 1, \hat{\eta}_{01} = \dots = \hat{\eta}_{0,b+1} = 1, \hat{\delta}_0 = 1$$

Also,  $\hat{\delta}_0 \geq \hat{\delta}_1 \geq \dots \geq \hat{\delta}_a$  are the eigenvalues of  $B$ ,  $\hat{\xi}_{u1}^*$  is the eigenvector of  $BB'$  corresponding to  $\hat{\delta}_u^2$  and  $\hat{\eta}_u^*$  is the eigenvector of  $B'B$  corresponding to  $\hat{\delta}_u^2$ . Now, let  $H_i: \delta_i^2 = 0$  ( $i = 1, 2, \dots, a$ ) and  $H = \bigcap_{i=1}^a H_i$ . We can use  $\psi(\hat{\delta}_1^2, \dots, \hat{\delta}_a^2)$  to test  $H$  where  $\psi(\cdot)$  is a suitable function of  $\hat{\delta}_1^2, \dots, \hat{\delta}_a^2$ . For example, we can use  $\hat{\delta}_1^2 + \dots + \hat{\delta}_a^2$  or  $\hat{\delta}_1^2$  as test statistics. Here we note that  $n(\hat{\delta}_1^2 + \dots + \hat{\delta}_a^2)$  is equivalent to  $\chi_0^2$  where

$$\chi_0^2 = \sum_{i,j} \{n_{ij} - (n_{i.} n_{.j}/n)\}^2 / n_{i.} n_{.j}. \quad (5.12)$$

When  $H_1$  is true, O'Neil [17] showed that the joint distribution of  $n\hat{\delta}_1^2, \dots, n\hat{\delta}_a^2$  is the same as the joint distribution of the eigenvalues of the central Wishart matrix with  $ab$  degrees of freedom. Percentage points of the largest eigenvalue of the central Wishart matrix are given in Krishnaiah [8].

We have discussed before some procedures to test for the overall hypothesis  $\delta_1^2 = \dots = \delta_a^2 = 0$ . We will now discuss procedures for testing the subhypotheses  $H_t$  when  $H_1$  is rejected. The hypothesis  $H_t$  is the same as the hypothesis that the rank of  $F$  is  $t$ . We will first consider the test procedure based upon  $T_1$  where  $T_q = \hat{\delta}_q^2 + \dots + \hat{\delta}_a^2$ . In this procedure, we accept or reject  $H_1$  according as

$$T_1 \leq c_{1\alpha} \quad (5.13)$$

where

$$P[T_1 \leq c_{1\alpha} | H_1] = (1 - \alpha). \quad (5.14)$$

If  $H_1$  is rejected, the hypothesis  $H_q$  is accepted or rejected according as

$$T_q \lesseqgtr c_{1\alpha}. \quad (5.15)$$

Starting with the test based upon  $T_1$  we can also draw inference on testing the hypothesis  $H_{ij}$  as described below; here  $H_{ij}$  denotes the hypothesis  $p_{ij} = p_i q_j$  for given values  $i, j$ . The  $\chi_0^2$  test statistic can be written as

$$\chi_0^2 = \underline{z}' \underline{z} \quad (5.16)$$

where

$$\underline{z}' = (z_{11}, \dots, z_{1,b+1}, \dots, z_{a+1,1}, \dots, z_{a+1,b+1})$$

and

$$z_{ij} = (n_{ij} - (n_{i.} n_{.j} / n)) / \sqrt{n_{i.} n_{.j}}.$$

But  $\chi_0^2 = \max(\underline{c}' \underline{z})^2$  when the maximum is taken over all non-null  $\underline{c}$  subject to the restriction that  $\underline{c}' \underline{c} = 1$ . So, when  $H_1$  is rejected, we can test the subhypotheses  $H_{ij}$  as follows. We accept or reject  $H_{ij}$  against two-sided alternatives according as

$$z_{ij}^2 \lesseqgtr c_{1\alpha}. \quad (5.17)$$

We can test the hypothesis  $\bigcap_{i=1}^t \bigcap_{j=1}^u H_{ij}$  as follows. We accept or reject

the above hypothesis against two-sided alternatives according as

$$\sum_{i=1}^t \sum_{j=1}^u z_{ij}^2 \lesseqgtr c_{1\alpha}$$

The hypothesis  $\bigcap_{j=1}^u \bigcap_{i=1}^t H_{ij}$  implies the hypothesis

$$\sum_{i=1}^t \sum_{j=1}^u p_{ij} = \left( \sum_{i=1}^t p_i \right) \left( \sum_{j=1}^u q_j \right).$$



We will now discuss the problem of testing the hypotheses  $H_{ij}$  against the alternatives  $A_{ij}$  simultaneously where  $A_{ij}: p_{ij} > p_{iq_j}$ . We accept or reject  $H_{ij}$  against  $A_{ij}$  according as

$$z_{ij} \leq c_{2\alpha} \quad (5.18)$$

where

$$P[\max z_{ij} \leq c_{2\alpha} | H_1] = (1-\alpha) \quad (5.19)$$

and  $\max z_{ij}$  denotes the maximum of the elements of  $\underline{z}$ . The asymptotic joint distribution of the elements of  $\underline{z}$  is a singular multivariate normal distribution. But, bounds on the critical values  $c_{2\alpha}$  can be obtained by using Poincare's formula. Similarly, we can propose procedures to test hypotheses  $H_{ij}$  against  $A_{ij}^*$  where  $A_{ij}^*: p_{ij} < p_{iq_j}$ .

We now discuss the test based upon  $\hat{\rho}_1^2$ . We accept or reject  $H_1$  according as

$$\hat{\rho}_1^2 \leq c_{2\alpha} \quad (5.20)$$

where

$$P[\hat{\rho}_1^2 \leq c_{2\alpha} | H_1] = (1-\alpha). \quad (5.21)$$

If  $H_1$  is rejected, we accept or reject  $H_j$  according as

$$\hat{\rho}_j^2 \leq c_{2\alpha}. \quad (5.22)$$

We will now derive the asymptotic nonnull distributions of certain functions of  $\hat{\rho}_1^2, \dots, \hat{\rho}_a^2$ . The following lemmas are needed in the sequel.

Lemma 5.1 Let  $U: p \times p$  be a symmetric matrix which can be expressed as

$$U = \Lambda + \epsilon U^{(1)} + \epsilon^2 U^{(2)} + \dots \quad (5.23)$$

when  $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_p)$ . Also, let

$$\lambda_{q_1 + \dots + q_{\alpha-1} + 1} = \dots = \lambda_{q_1 + \dots + q_{\alpha}} = \theta_{\alpha} \quad (5.24)$$

for  $\alpha = 1, 2, \dots, r$ ,  $q_0 = 0$  and  $q_r = a$ . In addition, let  $\lambda_1 \geq \dots \geq \lambda_p$  denote the eigenvalues of  $U$ . Then

$$\bar{\lambda}_{\alpha} = \theta_{\alpha} + \varepsilon \bar{\lambda}_{\alpha}^{(1)} + \varepsilon^2 \bar{\lambda}_{\alpha}^{(2)} + \dots \quad (5.25)$$

where

$$\begin{aligned} \bar{\lambda}^{(1)} &= \frac{1}{q_{\alpha}} \text{tr } U_{\alpha\alpha}^{(1)} \\ \bar{\lambda}^{(2)} &= \frac{1}{q_{\alpha}} \text{tr} \left[ U_{\alpha\alpha}^{(2)} + \sum_{\beta \neq \alpha} \theta_{\alpha\beta}^{-1} U_{\alpha\beta}^{(1)} U_{\beta\alpha}^{(1)} \right] \\ \theta_{\alpha\beta} &= \theta_{\alpha} - \theta_{\beta} \end{aligned}$$

$$U = \begin{bmatrix} U_{11}^{(i)} & U_{12}^{(i)} & \dots & U_{1r}^{(i)} \\ U_{21}^{(i)} & U_{22}^{(i)} & \dots & U_{2r}^{(i)} \\ \vdots & \vdots & & \vdots \\ U_{r1}^{(i)} & U_{r2}^{(i)} & \dots & U_{rr}^{(i)} \end{bmatrix}.$$

The above lemma is implicit in Kato [24]. It is also proved in Fujikoshi [5] by following the same lines as in Lawley [13].

**Lemma 5.2** Let  $\psi_i(\lambda_1, \dots, \lambda_a)$ , ( $i=1, 2, \dots, k$ ), be analytic functions of  $\lambda_1, \dots, \lambda_a$  around  $\lambda_1, \dots, \lambda_a$  and let  $\lambda_i$ 's have multiplicities as in (5.24). We assume that

$$\begin{aligned}
 \left. \frac{\partial T_1(\ell_1, \dots, \ell_a)}{\partial \ell_{j_1}} \right|_{\ell=\lambda} &= c_{1j_1} = a_{1\alpha} \\
 \left. \frac{\partial^2 T_1(\ell_1, \dots, \ell_a)}{\partial \ell_{j_1} \partial \ell_{j_2}} \right|_{\ell=\lambda} &= c_{1j_1 j_2} = a_{1\alpha\beta} \\
 \left. \frac{\partial^3 T_1(\ell_1, \dots, \ell_a)}{\partial \ell_{j_1} \partial \ell_{j_2} \partial \ell_{j_3}} \right|_{\ell=\lambda} &= c_{1j_1 j_2 j_3} = a_{1\alpha\beta\gamma}
 \end{aligned} \tag{5.26}$$

for  $j_1 \in J_\alpha$ ,  $j_2 \in J_\beta$ ,  $j_3 \in J_\gamma$ ,  $\ell' = (\ell_1, \dots, \ell_a)$ ,  $\lambda' = (\lambda_1, \dots, \lambda_a)$  and  $J_\alpha$  denotes the set of integers  $q_1 + \dots + q_{\alpha-1} + 1, \dots, q_1 + \dots + q_\alpha$ . Then

$$\begin{aligned}
 L_1 &= \sqrt{n} \{T_1(\ell_1, \dots, \ell_a) - T_1(\lambda_1, \dots, \lambda_a)\} \\
 &= \sum_{\alpha=1}^r a_{1\alpha} \text{tr} U_{\alpha\alpha}^{(1)} + \frac{1}{\sqrt{n}} \left[ \sum_{\alpha=1}^r a_{1\alpha} \text{tr}(U_{\alpha\alpha}^{(2)}) + \sum_{\beta \neq \alpha} \theta_{\alpha\beta}^{-1} U_{\alpha\beta}^{(1)} U_{\beta\alpha}^{(1)} \right. \\
 &\quad \left. + \frac{1}{2} \sum_{\alpha} \sum_{\beta} a_{1\alpha\beta} \text{tr} U_{\alpha\alpha}^{(1)} \text{tr} U_{\beta\beta}^{(1)} + \dots \right] \tag{5.27}
 \end{aligned}$$

for  $i = 1, 2, \dots, k$ . Let  $H$  and  $K$  be orthogonal matrices and let

$$R_1 = H' B K.$$

If we choose the first columns of  $H$  and  $K$  such that

$$h_{i0} = (n_{i.}/n)^{\frac{1}{2}} \quad i = 1, 2, \dots, (a+1)$$

$$k_{j0} = (n_{.j}/n)^{\frac{1}{2}} \quad j = 1, 2, \dots, (b+1),$$

then

$$R_1 = \begin{pmatrix} 1 & 0 \\ 0 & R \end{pmatrix}.$$

Similarly, let  $\Omega_1 = H^* F K^*$  where the first columns of  $H^*$  and  $K^*$  are given by

$$h_{i0}^* = p_i^{\frac{1}{2}}$$

$$k_{j0}^* = q_j^{\frac{1}{2}}$$

Then

$$\Omega_1 = \begin{pmatrix} 1 & 0 \\ 0 & \Omega \end{pmatrix}.$$

So,

$$R_1 R_1' = \begin{pmatrix} 1 & 0 \\ 0 & R R' \end{pmatrix}, \quad \Omega_1 \Omega_1' = \begin{pmatrix} 1 & 0 \\ 0 & \Omega \Omega' \end{pmatrix}.$$

It is known (see O'Neill [17]) that  $\hat{\rho}_1^2 \geq \dots \geq \hat{\rho}_a^2$  are the eigenvalues of  $R R'$  whereas  $\rho_1^2 \geq \dots \geq \rho_a^2$  are the eigenvalues of  $\Omega \Omega'$ . Next, let

$$X = \sqrt{n} (R - \Omega) = (x_{ij}) \quad (5.28)$$

where  $R = (r_{ij})$  and  $\Omega = (\omega_{ij})$ . Then  $x_{ij}$ 's are known (see O'Neill [17]) to be asymptotically distributed as multivariate normal with mean vector  $0$  and the elements of the covariance matrix are given by

$$\text{Cov}(x_{ij}, x_{kl}) = \sigma_{ij.kl} \text{ (say)}. \quad (5.29)$$

Using (5.28), we obtain

$$RR' = \Omega\Omega' + n^{-\frac{1}{2}}(\Omega X' + X\Omega') + n^{-1}XX'. \quad (5.30)$$

Now, let  $M: a \times a$  and  $L: b \times b$  be such that

$$M'\Omega L = (\text{diag}(\rho_1, \dots, \rho_a) | 0) = D_\rho.$$

Then,

$$S = M'RR'M = V + \frac{1}{\sqrt{n}} V^{(1)} + \frac{1}{n} V^{(2)} \quad (5.31)$$

where  $V = D_\rho D_\rho'$ ,  $V^{(1)} = (V_{\alpha\beta}^{(1)}) = D_\rho Z' + Z D_\rho'$ ,  $Z = (z_{ij}) = M'XL$ , and  $V^{(2)} = (V_{\alpha\beta}^{(2)}) = M'XX'M$ . Here  $V_{\alpha\beta}^{(1)}$  and  $V_{\alpha\beta}^{(2)}$  are of order  $q_\alpha \times q_\beta$ . Now, let  $\ell_i = \rho_i^2$ ,  $\lambda_i = \rho_i^2$  and  $\lambda_i$ 's have multiplicities as in (5.24). Then, using (5.26), we obtain

$$\begin{aligned} L_i &= \sqrt{n} \{T_i(\ell_1, \dots, \ell_p) - T_i(\lambda_1, \dots, \lambda_p)\} \\ &= \sum_{\alpha=1}^r a_{i\alpha} \text{tr } V_{\alpha\alpha}^{(1)} + \frac{1}{\sqrt{n}} \left[ \sum_{\alpha=1}^r a_{i\alpha} \text{tr} \{V_{\alpha\alpha}^{(2)} + \sum_{\beta \neq \alpha} \theta_{\alpha\beta}^{-1} V_{\alpha\beta}^{(1)} V_{\beta\alpha}^{(1)}\} \right. \\ &\quad \left. + \frac{1}{2} \sum_{\alpha, \beta} a_{i\alpha\beta} \text{tr } V_{\alpha\alpha}^{(1)} \text{tr } V_{\beta\beta}^{(1)} \right] \\ &\quad + \text{terms of higher order.} \end{aligned} \quad (5.32)$$

But,

$$V^{(1)} = \begin{pmatrix} 2\rho_1^2 z_{11} & (\rho_2^2 z_{12} + \rho_1^2 z_{21}) & \dots & (\rho_a^2 z_{1a} + \rho_1^2 z_{a1}) \\ (\rho_1^2 z_{21} + \rho_2^2 z_{12}) & 2\rho_2^2 z_{22} & \dots & (\rho_a^2 z_{2a} + \rho_2^2 z_{a2}) \\ \vdots & \vdots & \ddots & \vdots \\ (\rho_1^2 z_{a1} + \rho_a^2 z_{1a}) & (\rho_2^2 z_{a2} + \rho_a^2 z_{2a}) & \dots & 2\rho_a^2 z_{aa} \end{pmatrix} \quad (5.33)$$

$$V^{(2)} = \begin{pmatrix} M_1'XX'M_1 & M_1'XX'M_2 & \dots & M_1'XX'M_r \\ M_2'XX'M_1 & M_2'XX'M_2 & \dots & M_2'XX'M_r \\ \vdots & \vdots & \ddots & \vdots \\ M_r'XX'M_1 & M_r'XX'M_2 & \dots & M_r'XX'M_r \end{pmatrix}$$

where  $M = (M_1, \dots, M_r)$  and  $M_i$  is of order  $a \times q_i$ . So,

$$\begin{aligned} L_i &= 2 \sum_{\alpha=1}^r a_{i\alpha} \theta_{\alpha} (z_{q_1+\dots+q_{\alpha-1}+1, q_1+\dots+q_{\alpha-1}+1} + \dots + z_{q_1+\dots+q_{\alpha}, q_1+\dots+q_{\alpha}}) \\ &= \underline{b}_i' \underline{z}_0 \end{aligned} \quad (5.35)$$

where

$$\underline{b}_i' = (2a_{i1}\theta_1 1_{q_1}', \dots, 2a_{ir}\theta_r 1_{q_r}') \quad (5.36)$$

and  $\underline{z}_0' = (z_{11}, \dots, z_{aa})$ . The asymptotic distribution of  $B'\underline{z}$  is multivariate normal with mean vector  $\underline{0}$  and covariance matrix  $B'\Sigma^*B$  where  $\Sigma^*$  is the covariance matrix of  $\underline{z}$  and  $B = (\underline{b}_1, \dots, \underline{b}_k)$ . We can summarize the above results as follows:

Theorem 5.1 We assume that  $\rho_i$ 's have multiplicities as given below:

$$\rho_{q_1+\dots+q_{\alpha-1}} = \dots = \rho_{q_1+\dots+q_{\alpha}} = \theta_{\alpha} \quad (5.37)$$

for  $\alpha = 1, 2, \dots, r$  where  $q_0 = 0$ ,  $q_1 + \dots + q_r = a$ . Also, let  $L_1, \dots, L_k$  be functions of  $\hat{\rho}_1^2, \dots, \hat{\rho}_a^2$  satisfying the assumptions (5.25). Then, as  $n \rightarrow \infty$ , the joint distribution of  $L_1, \dots, L_k$  is multivariate normal with mean vector  $\underline{0}$  and covariance matrix  $B'\Sigma^*B$  where  $\Sigma^*$  is the covariance matrix of  $\underline{z}$ ,  $B = (\underline{b}_1, \dots, \underline{b}_k)$  and  $\underline{b}_i$ 's are defined by (5.36).

Now, let  $\sum_{\alpha=1}^r a_{i\alpha} \text{tr} V_{\alpha\alpha}^{(1)} = 0$  for each  $i$ . Then,

$$\begin{aligned} L_i^* &= n\{T_i(\ell_1, \dots, \ell_p) - T_i(\lambda_1, \dots, \lambda_p)\} \\ &= \frac{1}{2}(v_1, \dots, v_r) A_i \begin{pmatrix} v_1 \\ \vdots \\ v_r \end{pmatrix} \end{aligned}$$

where

$$\begin{aligned} A_i &= (a_{i\alpha\beta}) \\ v_1 &= 2(\rho_1 z_{11} + \dots + \rho_{q_1} z_{q_1 q_1}) \\ v_2 &= 2(\rho_{q_1+1} z_{q_1+1, q_1+1} + \dots + \rho_{q_1+q_2} z_{q_1+q_2, q_1+q_2}) \\ &\vdots \\ v_r &= 2(\rho_{q_1+\dots+q_{r-1}+1} z_{q_1+\dots+q_{r-1}+1, q_1+\dots+q_{r-1}+1} + \dots \\ &\quad + \rho_{q_1+\dots+q_r} z_{q_1+\dots+q_r, q_1+\dots+q_r}). \end{aligned}$$

Since  $\rho_i^2$ 's have multiplicities, we can write  $\underline{v} = (v_1, \dots, v_r)'$  as  $\underline{v} = E \underline{z}_0$  where

$$\begin{aligned} E &= (\underline{e}_1, \dots, \underline{e}_r), \\ \underline{e}_1' &= 2\theta_1(1'_{q_1}, 0, \dots, 0) \\ \underline{e}_2' &= 2\theta_2(0, \dots, 0, 1'_{q_2}, 0, \dots, 0) \\ &\vdots \\ \underline{e}_r' &= 2\theta_r(0, \dots, 0, 1'_{q_r}) \end{aligned}$$

As  $n \rightarrow \infty$ ,  $\underline{v}$  is distributed as a multivariate normal with mean vector  $\underline{0}$  and covariance matrix  $E \Sigma^* E'$ . So, the joint distribution of  $L_1^*, \dots, L_k^*$  is the same as that of correlated quadratic forms discussed in Khatrı, Krishnaiah and Sen [7].

# REFERENCES

- [1] Agresti, A. (1984). Analysis of Ordinal Categorical Data. John Wiley, New York.
- [2] Billingsley, P. (1968). Convergence of Probability Measures. John Wiley, New York.
- [3] Eaton, M.L. (1982). A review of selected topics in multivariate probability inequalities. Ann. Statist., 10, 11-43.
- [4] Fisher, R.A. and MacKenzie (1923). Studies in crop variation II. The manurial response of different potato varieties. J. Agric. Sci., 13, 311.
- [5] Fujikoshi, Y. (1978). Asymptotic expansions for the distributions of some functions of the latent roots of matrices in three situations. J. Multivariate Anal., 8, 63-72.
- [6] Goodman, L.A. (1985). The analysis of cross-classified data having ordered and/or unordered categories: association models, correlation models and asymmetry models for contingency tables with or without missing entries. Ann. Statist., 13, 10-69.
- [7] Khatri, C.G., Krishnaiah, P.R. and Sen, P.K. (1977). A note on the joint distribution of correlated quadratic forms. J. Statist. Plann. Inference, 1, 299-307.
- [8] Krishnaiah, P.R. (1980). Computations of some multivariate distributions. In Handbook of Statistics, Vol. 1: Analysis of Variance, pp. 745-971. North-Holland Publishing Company.
- [9] Krishnaiah, P.R. and Lee, J.C. (1979). On the asymptotic joint distributions of certain functions of the eigenvalues of some random matrices. J. Multivariate Anal., 9, 248-258.
- [10] Krishnaiah, P.R. and Yochmowitz, M. (1980). Inference on the structure of interaction in two-way classification model. In Handbook of Statistics, Vol. 1: Analysis of Variance, pp. 973-994. North-Holland .
- [11] Lancaster, H.O. (1958). The structure of bivariate distributions, Ann. Math. Stat., 29, 719-736.
- [12] Lancaster, H.O. (1969). The Chi-Squared Distribution. John Wiley, New York.
- [13] Lawley, D.N. (1956). Tests of significance for the latent roots of covariance and correlation matrices. Biometrika, 43, 128-136.
- [14] Lehmann, E.L. (1966). Some concepts of dependence. Ann. Math. Stat., 37, 1137-1153.



- [15] Mandel, J. (1969). Partitioning the interaction in analysis of variance. J. Res. Nat. Bur. Standards, Sect. B, 73B, 309-328.
- [16] Nguyen, T.T. and Sampson, A.R. (1983). Testing for positive quadrant dependence in ordinal contingency tables. Technical Report No. 83-10, Center for Multivariate Analysis, University of Pittsburgh, Pittsburgh.
- [17] O'Neill, M.E. (1978). Asymptotic distributions of the canonical correlations from contingency tables, Austral. J. Statist., 20, 75-82.
- [18] O'Neill, M.E. (1978). Distributional expansions for canonical correlations from contingency tables, J. Roy. Statist. Soc. Ser. B, 40, 303-312.
- [19] O'Neill, M.E. (1980). A note on canonical correlations from contingency tables. Austral. J. Statist.,
- [20] Phelps, R.R. (1966). Lectures on Choquet's Theorem. Van Nostrand, New York.
- [21] Subramanyam, K. and Bhaskara Rao, M. Extreme point methods in the study of classes of bivariate distributions and some applications to contingency tables. To be issued as a Technical Report.
- [22] Tukey, J.W. (1949). One degree of freedom for non-additivity. Biometrics, 5, 232-242.
- [23] Williams, E.J. (1952). The interpretation of interactions in factorial experiments. Biometrika, 39, 65-81.
- [24] Kato, T. (1976). Perturbation Theory for Linear Operators. Springer Verlag, New York.
- [25] Goodman, L.A. and Kruskal, W.H. (1979). Measures of Association for Cross Classification. Springer Verlag, New York.

END

DT/C

8-86